

## **AGGRESSIVE SURGICAL TREATMENT IN CERVICAL TUMOR WITH MULTIPLE LOWER ABDOMINAL DISSEMINATIONS**

Irina Bălescu<sup>1</sup>, N. Bacalbașa<sup>2</sup>, Claudia Stoica<sup>3</sup>, R. Sgarbura<sup>3</sup>, Dana Tomescu<sup>4</sup>

<sup>1</sup>“Ponderas” Hospital, Bucharest, Romania

<sup>2</sup>“Carol Davila” University of Medicine and Pharmacy, Bucharest, Romania

<sup>3</sup>“Ilfov” County Hospital, Bucharest, Romania

<sup>4</sup>Department of Anesthesia and Intensive Care III, Fundeni Clinical Institute Bucharest Romania

### **Abstract**

In cases with advanced cervical cases, with large lymph node metastases or solitary peritoneal masses the patient is candidate rather for debulking surgery than systemic treatments. Several studies have proven that radiotherapy is inefficient for tumoral masses larger than 2 cm. In this case report we present a 55 years old patient with advanced cervical cancer, large lymph node metastasis and a solitary peritoneal mass. A radical hysterectomy with bilateral adnexectomy, lymph node dissection and mesosigmoid mass resection were performed. The patient was addressed to adjuvant chemotherapy.

**Key words:** advanced cervical cancer, large lymph node metastases, debulking surgery, irradiation

Address for correspondence: Nicolae Bacalbașa, MD  
Dimitrie Racoviță Street, no. 2, Bucharest, Romania  
E-mail: nicolae\_bacalbasa@yahoo.ro